

Article

# Olfactory Characterization of Typical Odorous Pollutants Part I: Relationship Between the Hedonic Tone and Odor Concentration

Jiayin Li<sup>1,2</sup>, Kehua Zou<sup>1,3</sup>, Weifang Li<sup>1,3,\*</sup>, Gen Wang<sup>1,3</sup> and Weihua Yang<sup>1,2</sup>

- <sup>1</sup> State Environmental Protection Key Laboratory of Odor Pollution Control, Tianjin 300191, China
- <sup>2</sup> Tianjin Sinodour Environmental Technology Co., Ltd., Tianjin 300191, China
- <sup>3</sup> Tianjin Academy of Environmental Sciences, Tianjin 300191, China
- \* Correspondence: lwf1919@163.com

Received: 29 July 2019; Accepted: 3 September 2019; Published: 6 September 2019



**Abstract:** The hedonic tone is a suitable evaluation index which can truly reflect the psychological impact of an odor. To find out the relationship between the odor concentration (OC) and hedonic tone (H), dimethyl disulfide, limonene and butyl acetate were presented as typical odorants with different characters. A panel of 16 persons was engaged to rate the hedonic tone of a series sample with various concentrations according to the nine-point scale. The relationship between the hedonic tone and OC was established based on a multivariate logistic regression analysis. The research results demonstrated that the smell of dimethyl disulfide is unpleasant at various concentration levels, and its perceived unpleasantness is increased with OC, and at the critical point (H = -0.5), the odor index of dimethyl disulfide is 0.5 (OC =  $3 \text{ OU}_{\text{E}} \cdot \text{m}^{-3}$ ). For limonene, its smell is pleasant when the odor index is between 1.4 and 3.3 (OC =  $25 \sim 1995 \text{ OU}_{\text{E}} \cdot \text{m}^{-3}$ ). For butyl acetate, the average results showed an unpleasant character with the corresponding odor index of 1.87 (OC =  $74 \text{ OU}_{\text{E}} \cdot \text{m}^{-3}$ ). Each odorant has a unique hedonic behavior curve from which the annoyance potential of different odorants can be clearly discriminated, with the order of dimethyl disulfide > butyl acetate > limonene. The regression equations showed a quadratic nonlinear function between the hedonic tone and OC.

**Keywords:** typical odorants; hedonic tone; odor concentration; hedonic curve; quantitative relationship

# 1. Introduction

Odor is an important component of air pollution that can cause people annoyance by irritating the nose and lead to a negative quality of life [1,2]. Although the ambient air quality has significantly improved in the last few decades, the tolerance of residents to odor impacts appears to be reduced [3]. It is difficult to quantitatively analysis the odor substances because of their low olfactory threshold concentration and high chemical activity [4]. In addition, odor pollution is based on the psychological impact of people as the main features [5], for example: Vanillin with a fragrant scent will make people feel pleasant, while Ammonia with a pungent smell will make people feel disgusted, but the gas composition concentration analyzed by the instrument cannot truly reflect the sensory impact on the human organ [6]. Therefore, odor is a sensation measurable through the subjective responses of individuals. The extent of the annoyance caused by odorous air depends on a number of parameters, collectively known as FIDOL, namely the frequency (F), intensity (I), duration (D), offensiveness (O), and location (L) [7–9]. The offensiveness of an odor perception is often measured in terms of the "hedonic tone", a category judgment of the relative like (pleasantness) or dislike (unpleasantness) of the odor [10]. Furthermore, experience has shown that the hedonic tone can reflect injury and the



psychological influence better than the most widely used regulatory parameter—odor concentration, OC [11].

The odors emitted by various industrial sources differ by their nuisance potential because the hedonic tone of the odor apparently affects the degree of the nuisance [12,13]. Winneke and Kastka [14] reported the importance of the hedonic tone on human beings' psychological feelings. They surveyed residents near a candy factory, an insulation plant, a tar refinery and a winery, and found that the annoyance caused by different industrial sources was not the same. The chocolate factory is rated as being less unpleasant and harassing than the tar refinery. A similar test was conducted by Hangartner and Wüst, and the results showed that the odors from grass-drying plants and food-processing plants caused less annoyance than those in the chemical industry and in tar processing [15]. In a research about odor annoyance evaluation, the prediction of the percentage of those being "highly annoyed" (HA%) would be much closer to the actual situation if the hedonic tone was taken into account. The HA% at a certain exposure level will be lower than the more unpleasant odor [16]. A related site investigation was conducted around six odor emitting sources; two were pleasant, two were neutral, and two were unpleasant odor emissions, the results showing that the hedonic tone is a powerful predictor of annoyance, and that the exposure-annoyance as well as exposure-symptom associations are strongly affected by the hedonic tone [17]. In addition, the hedonic tone is an important dimension for the odor classification by humans with diverse cultural backgrounds [18]. The same conclusion was obtained by Wang et al. [19], in her research about 60 odor words that were classified by ninety college students based on the individual perceptual similarity; the results showed that the hedonic tone can be the standard of odor words classification.

Currently, the control and regulation of odor pollution is mostly based on the OC parameter [20]. The OC is defined as being the dilution factor required to reach the sensory threshold. From the definition, we can see that the OC is independent of the odor character itself, and therefore cannot truly reflect the personal perception of diverse odors [21]. For example, the smell of perfume and sewage may have the same OC, but their psychological feelings are completely different, and the acceptable concentration levels are inevitably different. Practical experience has shown that the OC is not an adequate criterion for the evaluation of a variety of odors. Additional factors, such as their hedonic tone, must also be taken into account [22,23]. In some countries (e.g., Germany, Ireland, and Belgium), the odor impact criteria can be adapted according to the hedonic tone of an odor by using a weighting factor [24].

For a certain odorant or odorous sample, its hedonic tone is not fixed but varies with the concentration. Understanding the relation between the hedonic tone and OC is helpful for the determination of acceptable concentration levels as well as for setting odor impact criteria. Nimmermark [25] conducted research to study the influence of the odor concentration on the hedonic tone of odor from livestock and poultry farm. The results showed that at the same hedonic level of -1, which means slightly unpleasant, the corresponding odor concentration is 14–16 OUE·m<sup>-3</sup> from livestock, while it is 37 OUE·m<sup>-3</sup> from a cow shed. Huang and Guo [26] conducted research in a commercial daily barn, layer barn, and broiler barn; a reference table of OC limits was generated with 3 set values for the hedonic tone (0, -1, and -2) for all three odor sources, which may provide references in establishing appropriate odor impact criteria to meet different land use purposes.

As yet, there have been few studies about the hedonic tone in China. The main purpose of this article is to present the hedonic characteristics of three typical odorous substances: dimethyl disulfide, limonene and butyl acetate, which are widely distributed in the environment. Additionally, the purpose of this article is improve the odor evaluation criterion in China by studying the relationship between the hedonic tone and OC. The achievements can lay the foundation for a more scientific odor pollution characterization method.

# 2. Experimental Section

#### 2.1. Standard Substances

In the research, more than 20 odorants were tested. Furthermore, three of them, dimethyl disulfide, butyl acetate and limonene, were chosen as the representative odorants with different hedonic characteristics. Dimethyl disulfide is a flammable liquid with an unpleasant, garlic-like odor. Limonene is a colorless liquid hydrocarbon that possesses a pleasant smell of oranges. Butyl acetate is found in many types of fruit; however, there is a divergence about the smell: some people feel that it is pleasant, like a sweet smell of banana or apple, but for others it is unpleasant. In this research, 50 ppm of dimethyl disulfide and limonene gas and 500 ppm of butyl acetate gas were selected as the original samples.

### 2.2. Sample Preparation

The standard gas was diluted to 6 concentration gradients that differ by a factor of approximately 3. The lower limit must correspond to the panel threshold. The higher limit must be checked for toxicity, and all possible health risk of the panelists must be totally excluded. The dilutions were filled into the 8 L polyester odorless sampling bags respectively as the testing sample, and inserted into two bags of nitrogen as the blank samples. The process was done with the help of a dynamic dilution apparatus (4600, Entsch, USA).

# 2.3. Testing Method

The OC of a sample is measured by diluting the sample with odorless air until it cannot be discriminated from odorless air by a test group of persons. The OC of the original standard gaseous samples were determined following the guideline "Air Quality-Determination of Odor-Triangle Odor Bag Method" [27] The determination of the hedonic tone refers to a 9 point scale with values ranging from "-4-extremely unpleasant" to "+4-extremely pleasant", via "0-neither pleasant nor unpleasant" ("neutral") (See Table 1).

Hedonic Tone	Verbal Description
-4	extremely unpleasant
-3	moderate unpleasant
-2	unpleasant
-1	slightly unpleasant
0	neutral
1	slightly pleasant
2	pleasant
3	moderate pleasant
4	extremely pleasant

Table 1. The 9-point hedonic scale for odors.

In order to prevent the interference factors, it should be ensured that the experiment room is odorless and has good ventilation. The temperature of all samples is 26~28 °C, which is comfortable for the human body.

The presentation of the odor samples with different concentrations was done at random, but the series of measurements must start neither with a blank sample nor with a maximum concentration. In order to avoid panelists' possible guessing tendencies, the presentations were interspersed with blank ones.

The following instructions were given to the assessors before the test: The hedonic tone of an odor must not be confused with its quality (fragrant, pungent, stale or smelling like ... ). There is no "right" or "wrong" answer. The results are just on behalf of the personal feeling. The panel hedonic tone is the average value of all the panelists.

#### 2.4. Panel of Assessors

In order to avoid the differences between individuals that may occur when determining the hedonic odor tone, the panel consisted of 16 qualified members who have passed the Sniffer Certification Test according to the Air Quality-Determination of Odor-Triangle Odor Bag Method. Seven of them were female and nine were male. Their ages varied from 22 up to 45, with a mean of 35 years. All the panelists came from different cities from the south to the north of China, with different living habits.

The panelist needs to be normalized by a 5-day routine course to make sure that the data are standard, accurate and stabilized. Each day, they were all trained with two reference materials: vanillin (5 g/L, dipropylene glycol) and guaiacol (5  $\mu$ L/L, distilled water). The assessors rated the hedonic tone according to the 9-point hedonic scale. Before the test, they were instructed that "extremely unpleasant" should represent the most unpleasant odor they had experienced and that "extremely pleasant" should represent the most pleasant odor they had experienced. Their results were recorded every day. According to the requirements of Verein Deutscher Ingenieure, VDI 3882, the mean value of the results obtained from the entire panel must lie within a given range as follows:

- 1. Vanillin: from +2.9 to +1.9
- 2. Guaiacol: from -0.8 to -2.0

All the study participants gave written informed consent. The study was approved by the local research ethics committee.

## 2.5. Data Processing Method

The data fitting module of Origin was used to analyse the relationship between the hedonic tone and OC. Based on the human psychological response to odors, the OCs were logarithm transformed, which is referred to as the odor index:

$$C' = \log C \tag{1}$$

where C' is the odor index, and C is the OC.

#### 3. Results

#### 3.1. The Hedonic Tone of Dimethyl Disulfide

Dimethyl disulfide is a typical odorant. According to the "Air Quality-Determination of Odor-Triangle Odor Bag Method", the original OCs of the sample were measured as the result of 7413  $OU_E \cdot m^{-3}$ . The original sample was diluted to 10, 30, 100, 300, 1000 and 3000 times, which were represented by  $Z_1 \sim Z_6$ . The hedonic tones at different OCs were determined by the panel members, and the results are shown in Table 2.

Generally, the dimethyl disulfide hedonic tones were all negative, which means that they belong to the range of unpleasant substances. For the concentration level of  $Z_1$ , about 56% of panelists gave the answer "extremely unpleasant", and 38% "moderately unpleasant". As the dilution factor increases, the degree of aversion gradually mitigates. Until the concentration was diluted to  $Z_6$ , the majority was "neither pleasant nor unpleasant", and others were "slightly unpleasant". The average values of the hedonic tone from  $Z_1$  to  $Z_6$  are: -3.50, -2.75, -1.69, -1.56, -0.81, and -0.38.

In addition, hydrogen sulfide, ammonia, methyl mercaptan and some other odorants are similar to dimethyl disulfide. The odors at each dilution factor are all unpleasant, and the aversion gradually diminishes as the dilution multiples increase. The substances with such hedonic characteristics are all called unpleasant odorants.

Hedonic	Z <sub>1</sub>		Z <sub>2</sub>		Z <sub>3</sub>		$Z_4$		Z <sub>5</sub>		Z <sub>6</sub>		Blank Sample A		Blank Sample B	
Tone	Q	Р	Q	Р	Q	Р	Q	Р	Q	Р	Q	Р	Q	Р	Q	Р
4	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%
3	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%
2	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%
1	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	1	6%	0	0%
0	0	0%	0	0%	0	0%	0	0%	4	25%	10	62%	15	94%	16	100%
-1	0	0%	1	6%	6	38%	8	50%	11	69%	6	38%	0	0%	0	0%
-2	1	6%	4	25%	9	56%	7	44%	1	6%	0	0%	0	0%	0	0%
-3	6	38%	9	56%	1	6%	1	6%	0	0%	0	0%	0	0%	0	0%
-4	9	56%	2	13%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%
Average value	-3	3.50	-	2.75	-	1.69		1.56	-(	).81	-0	0.38	0.	.06	(	).00

**Table 2.** The result of the hedonic tone evaluation for dimethyl disulfide; the odor concentration, OC of the original sample is 7413  $OU_E \cdot m^{-3}$ . The dilution times from  $Z_1$  to  $Z_6$  are respectively 10, 30, 100, 300, 1000 and 3000. Q: amount of people; P: proportion of the population.

#### 3.2. The Hedonic Tone of Limonene

1

0

 $^{-1}$ 

-2

-3

-4

Average

value

0

0

3

3

6

2

0%

0%

19%

19%

37% 0

12%

-1.81

1

0

3

4

0

6%

0%

19%

25%

0%

0%

0.56

5

0

2

2

0 0%

0

31%

0%

12%

12%

0%

0.81

12

1

1

0

0

0

Limonene is an odorant with the scent of citrus. The original OC is 7413  $OU_E \cdot m^{-3}$ . Although the concentration was the same as dimethyl disulfide, the odor characteristics were completely different. The smell of limonene was not as strong as dimethyl disulfide, and was less unbearable. The concentration gradients include: the original gas, 10, 30, 100, 300 and 1000 dilution steps. The determination results are shown in Table 3.

OU <sub>E</sub> ·m	-3.7	The dil	utio	n time	s fro	$m Z_1 t$	o Z <sub>6</sub> a	are resp	ectiv	ely 0, 1	0, 30,	100, 300	and 1	000.		
Hedonic	:	Z1		Z <sub>2</sub>		Z <sub>3</sub>	2	$\mathbb{Z}_4$	7	<b>Z</b> <sub>5</sub>	2	Z <sub>6</sub>	Bl Sam	lank 1ple A	Bl San	ank 1ple B
Tone	Q	Р	Q	Р	Q	Р	Q	Р	Q	Р	Q	Р	Q	Р	Q	Р
4	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%
3	2	13%	3	19%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%
2	0	0%	5	31%	7	45%	2	12%	0	0%	0	0%	0	0%	0	0%

76%

6%

6%

0%

0%

0%

0.94

1

15

0

0

0

0

0.06

6%

94%

0%

6%

0%

0%

0

16

0

0

0

0

0%

100%

0%

0%

0%

0%

0.00

0

16

0

0

0

0

0%

100%

0%

0%

0%

0%

0.00

0

16

0

0

0

0

0%

100%

0%

0%

0%

0%

0.00

**Table 3.** The result of the hedonic tone evaluation for limonene; the OC of the original sample is 7413  $OU_E \cdot m^{-3}$ . The dilution times from  $Z_1$  to  $Z_6$  are respectively 0, 10, 30, 100, 300 and 1000.

The smell of limonene varied with its OC. For the undiluted standard gas, the average value of the hedonic tone was –1.81, which was close to "moderately unpleasant." When diluted 10 times, there were nine panelists who thought the smell was pleasant, seven felt it was unpleasant, and the average hedonic tone was 0.56, which was between "slightly pleasant" and "neither pleasant nor unpleasant." When diluted 30 times, there were 12 panelists who thought that it was pleasant, and the average hedonic tone was 0.80, which was close to "slightly pleasant." When diluted 100 times, 14 panelists thought it was pleasant, and the average hedonic tone almost reached 1, meaning "slightly pleasant." When diluted 300 times, the odor became very weak, about 94% of the panelists thought it was "neither pleasant nor unpleasant," and the average value of the hedonic tone was near 0. When diluted 1000 times, none of the panelists could perceive anything. This indicated that the smell of limonene usually provides a pleasant feeling, but that the odor decays relatively fast.

Besides, ethyl acetate, vanillin and some other scented substances have the similar hedonic characteristics to limonene. The odors are pleasant at lower concentration gradients, while they are unpleasant at higher concentration gradients. The substances with such hedonic characteristics are called pleasant odorants.

# 3.3. The Hedonic Tone of Butyl Acetate

The OC of butyl acetate is 17,378  $OU_E \cdot m^{-3}$ , and the serial samples of the test include: the original gas, 10, 30, 100, 300 and 1000 dilution steps. The results of the evaluation from the test group are shown in Table 4.

**Table 4.** The result of the hedonic tone evaluation for butyl acetate; the OC of the original sample is 17,378  $OU_E \cdot m^{-3}$ . The dilution times from  $Z_1$  to  $Z_6$  are respectively 0, 10, 30, 100, 300 and 1000.

Hedonic	:	Z <sub>1</sub>		Z <sub>2</sub>		Z <sub>3</sub>		$Z_4$		$Z_5$	2	Z <sub>6</sub>	B San	lank nple A	B San	lank nple B
Tone	Q	Р	Q	Р	Q	Р	Q	Р	Q	Р	Q	Р	Q	Р	Q	Р
4	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%
3	2	13%	1	6%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%
2	0	0%	1	6%	2	13%	1	6%	0	0%	0	0%	0	0%	0	0%
1	0	0%	0	0%	0	0%	3	19%	5	31%	1	6%	0	0%	0	0%
0	0	0%	0	0%	0	0%	1	6%	2	13%	11	69%	16	100%	16	100%
-1	2	13%	2	13%	5	31%	7	44%	6	37%	4	25%	0	0%	0	0%
-2	0	0%	4	25%	4	25%	4	25%	2	13%	0	0%	0	0%	0	0%
-3	2	13%	5	31%	5	31%	0	0%	1	6%	0	0%	0	0%	0	0%
-4	12	75%	3	19%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%
Average value	-(	3.13	_	2.00	_	1.50	_	0.63	_	0.50	-(	).19	(	).00	0	).00

In general, the smells of butyl acetate at different OC steps were all unpleasant, and the average values from  $Z_1$  to  $Z_6$  were respectively -3.12, -2.00, -1.50, -0.63, -0.50, and -0.19. The hedonic tone of butyl acetate decreases as its concentration decreases. However, different from disulfide, there were significant distinctions among the individuals for butyl acetate, insofar as when it was diluted 10 times, 2 panelists thought that it was pleasant. The number of pleased panelists increased with the dilution multiples, and when the odor concentration was diluted 300 times, about 5 people thought it was pleasant.

Methylisobutanone and propionaldehyde are similar to butyl acetate. At a same odor concentration, some people find this pleasant and some people find this unpleasant. Substances with such hedonic characteristics are called divergent odorants.

# 4. Discussion

# 4.1. The Relationship Between the Hedonic Tone and OC

The behavior curves of each odorant were plotted in the study with the odor index as the abscissa and the hedonic tone as the ordinate. The regression equations of the hedonic tone and the odor index were fitted through the data analysis software of Origin. According to these characteristic curves, the hedonic tone at different OCs can be obtained.

Figure 1 shows the relationship between the hedonic tone of dimethyl disulfide and the odor index. When the odor index < 0.5 (OC =  $3 \text{ OU}_{\text{E}} \cdot \text{m}^{-3}$ ), the average value of the hedonic tone is similar to 0, indicating that panel members feel neither pleasure nor displeasure. Therefore, the OC  $3 \text{ OU}_{\text{E}} \cdot \text{m}^{-3}$  can be used as the environmental impact reference value for dimethyl disulfide. Although the odor becomes more and more repulsive with the increase of the odor index, the relationship between them

is not linear. There is a non-linear binomial relationship between the hedonic tone and the odor index; the correlation coefficient is 0.97, and the fitting model is:



$$Y = -0.22 - 0.45X - 0.25X^2 (R^2 = 0.97)$$
(2)

**Figure 1.** The behavior curve of dimethyl disulfide. The black line is the actual line, the red line is the fitting line and the red point is the hedonic tone of -0.5.

However, the hedonic tone is not only related to the OC but also to various complex factors, such as the odor intensity, odor properties, and personal life experience [28]. Therefore, the formula can only be used to reflect the incremental relationship between the OC and hedonic tone, but not for an accurate prediction of the hedonic tone.

Figure 2 shows the relationship between the hedonic tone of limonene and the odor index. One can see the change process of the hedonic tone from pleasant to repulsive as the concentration increases. The panel members feel pleasure when the odor index is between 1.4 and 3.3 ( $OC = 25 \sim 1995 OU_E \cdot m^{-3}$ ). Additionally, the hedonic value reaches the maximum as the concentration index reaches 2.0 ( $OC = 100 OU_E \cdot m^{-3}$ ), after which point it starts to decline. If the odor index exceeds 3.3 ( $OC = 1995 OU_E \cdot m^{-3}$ ), the smell will change to unpleasant. 1995  $OU_E \cdot m^{-3}$  can be used as a reference value for the environmental impact of limonene. The hedonic curve of limonene clearly shows that when the concentration of an odorant exceeds a certain value, its smell becomes repulsive, even if it smells pleasant at low or general concentration levels. Through data fitting, a binomial relationship between the hedonic tone and odor index is obtained with a correlation coefficient of 0.99, and the relationship model is:

$$Y = -2.33 + 3.28X - 0.81X^2 (R^2 = 0.99)$$
(3)

Figure 3 shows the behavior curve of butyl acetate. Similar to dimethyl disulfide, the hedonic tones of butyl acetate are all negative values in the testing concentration range, and the degree of unpleasantness is enhanced with the increase of the odor index. When the odor index < 1.87 ( $OC = 74 \text{ OU}_{\text{E}} \cdot \text{m}^{-3}$ ), the hedonic tone is approximately 0, and the smell of butyl acetate is neutral, which will not make people get annoyed. Thus, 74  $OU_{\text{E}} \cdot \text{m}^{-3}$  can be used as the environmental impact reference value for butyl acetate. There was a binomial relationship between the hedonic tone and the odor index, and the correlation coefficient was 0.99. The relationship model is:

$$Y = -0.13 + 0.30X - 0.27X^2 (R^2 = 0.99)$$
(4)



Figure 2. The behavior curve of limonene.



Figure 3. The behavior curve of butyl acetate.

#### 4.2. Comparison of the Hedonic Curves of the Three Odorants

The hedonic curves of dimethyl disulfide, limonene and butyl acetate are plotted in a graph (see Figure 4), which clearly shows the characteristics and differences. The curves of dimethyl disulfide and butyl acetate all fell in the "unpleasant" region, but the degree of unpleasantness of dimethyl disulfide was higher than that of butyl acetate for the same odor index. For example, when the odor index is 2 (OC = 100 OU<sub>E</sub>·m<sup>-3</sup>), the hedonic tone of butyl acetate is -0.6, while the value of dimethyl disulfide is -2.0. The change rate of the hedonic tone with the odor index is different for the three substances, in the order of dimethyl disulfide > butyl acetate > limonene. The smell of dimethyl disulfide has a greater impact on the population.

The perceived smell is pleasant when the hedonic tone exceed +0.5, while it is unpleasant when the value is less than -0.5. The OC for the hedonic tones of +0.5 and -0.5 is defined as the critical point at which the smell of an odorant will change to pleasant or unpleasant. For dimethyl disulfide and butyl acetate, the unpleasant critical point is  $3 \text{ OU}_{\text{E}} \cdot \text{m}^{-3}$  and  $74 \text{ OU}_{\text{E}} \cdot \text{m}^{-3}$ . What is different with them is that limonene has two critical points:  $25 \text{ OU}_{\text{E}} \cdot \text{m}^{-3}$  is the pleasant point and 1995  $\text{OU}_{\text{E}} \cdot \text{m}^{-3}$ is the unpleasant point. The facts show that the OC above the unpleasant critical points will more likely cause an environment interference, which may cause adverse effects on humans, like annoyance, nuisance and complaint.



Figure 4. Comparison of the behavior curves of the three typical odorants.

The emission standard of odor pollution in China is mainly in relation to the OC. However, odor is a sensory pollution, with a strong subjectivity, and different odorants may have different critical points for causing unpleasant feelings. The practices at home and abroad have confirmed that the OC limits, without considering the hedonic tone, may be excessive and inadequate. It is therefore important for environmental management to develop different odorants' unpleasant critical points to improve the existing standards. The hedonic curves provide a scientific basis for the determination of the limit.

#### 4.3. The Panelist Response Curves

The hedonic curve is plotted according to the average value of the whole panel, and it can reflect the characteristics of the overall change. The personal response curve can be drawn through the result of each panelist. The data showed that the panelist response curves of butyl acetate are in the "unpleasant" region but that there is a significant difference between the individuals. 4 members' individual results for butyl acetate are shown in Figure 5. One can see a process from pleasant to unpleasant as the OC increases. Furthermore, the maximum and the turning point of the hedonic tone are different for different members.



Figure 5. The behavior curves of individual members for butyl acetate.

There are many reasons for individual differences, i.e., age, gender, living environment, and so on. Table 5 shows the unpleasant critical points of different ages and genders. For dimethyl disulfide, the panelists with different human factors have similar results of  $3 \text{ OU}_{\text{E}} \cdot \text{m}^{-3}$ . This is because the odor threshold of dimethyl disulfide is very low, and people feel unpleasant once receiving it. For limonene and butyl acetate, the unpleasant critical points for females are lower than those for males. This suggests that females can be aware of odors in the environment when concentrations are

low. In a number of studies regarding odor detection, women have performed better than men [29], and it has therefore also been suggested that women might complain more about unpleasant odors in the environment than men [30].

Charac	teristic	Average Value ( $OU_E \cdot m^{-3}$ )	SD	Odds, 95% CI	CV							
Dimethyl Disulfide												
Gender	Female	2	0.55	1.65-2.35	0.28							
	Male	4	1.22	3.33-4.67	0.31							
Ago	22~35	3	1.41	2.28-3.72	0.47							
Age	35~45	3	1.11	2.43-3.57	0.37							
Limonene												
Gender	Female	1836	124.92	1757.20-1914.80	0.07							
	Male	2119	203.02	2008.02-2229.98	0.10							
Age	22~35	1915	185.84	1819.76-2010.24	0.10							
	35~45	2128	216.52	1980.04-2275.96	0.10							
Butyl acetate												
Gender	Female	50	15.52	40.21-59.79	0.31							
	Male	39	14.26	31.20-46.80	0.37							
Age	22~35	65	26.23	51.56-78.44	0.40							
	35~45	89	18.64	76.26-101.74	0.21							

Table 5. The unpleasant critical points of different genders and ages.

Furthermore, the age of the panelists can influence the results. In this research, the unpleasant critical points of the panelists aged 22 to 35 are generally lower than those aged 35 to 45. This is because young people's noses are more sensitive than those of the elder. At the same odor concentration, the persons with sensitive noses are likely to find more unpleasantness than those with insensitive noses.

In addition to the factors of age and gender, some extra characteristics may also influence the results, like the living habit, place of residence (countryside or village), profession, and so on. This is the reason why the determination of the hedonic tone should consist of at least 15 persons.

# 5. Conclusions

The characteristics of the hedonic tone vary for different odor types. Dimethyl disulfide is an unpleasant odorant, and the degree of disgust gradually increases with the OC. Limonene belongs to the category of pleasantness for a certain OC, and it will change to disgust if the OC is beyond that range. Butyl acetate is unpleasant in general, but there divergences exist among individuals. The mathematical equations developed using the multivariate regression analysis show a quadratic nonlinear relationship between the hedonic tone and OC. The annoyance potential that was sorted according to the hedonic curves is: dimethyl disulfide > butyl acetate > limonene. In general, there is only one critical point for odorants with an unpleasant smell, while odorants with a pleasant smell usually present two critical points: a pleasant one and an unpleasant one.

However, the critical points are theoretical values obtained through the hedonic curves. Research has shown that age and gender factors may influence the results. In our future work, the results will be verified in practical application. Furthermore, studies about the hedonic characteristics of odor pollution sources, i.e., wastewater treatment plants, landfills, and livestock farms, will be carried out in the next step.

Author Contributions: Conceptualization, J.L. and W.L.; methodology, J.L. and W.Y.; validation, W.L. and K.Z.; investigation, J.L. and W.Y.; resources, K.Z. and G.W.; writing—original draft preparation, J.L.; writing—review and editing, K.Z. and W.L.; visualization, K.Z. and G.W.; supervision, W.L.

**Funding:** This research was funded by NATIONAL NATURAL SCIENCE FOUNDATION OF CHINA, grant number 21577096.

**Acknowledgments:** I would like to show my deepest gratitude to Gunther and Martin who have provided me with valuable guidance in the writing of this thesis. I shall extend my thanks to all my colleagues for their help in the experiment.

**Conflicts of Interest:** The authors declare no conflict of interest.

# References

- Donham, K.J. Community and occupational health concerns in pork production: A review. J. Anim. Sci. 2010, 88, 102–111. [CrossRef]
- 2. Wing, S.; Horton, R.A.; Marshall, S.W.; Thu, K.; Taiik, M.; Schinasi, L. Air pollution and odor in communities near industrial swine operations. *Environ. Health Persp.* **2008**, *116*, 136–1368. [CrossRef]
- 3. Guffanti, P.; Pifferi, V.; Falciola, L.; Ferrante, V. Analyses of odours from concentrated animal feeding operations: A review. *Atmos. Environ.* **2018**, *175*, 100–108. [CrossRef]
- 4. Jacobsen-Garcia, A.; Dale, M.; Desrochers, R.; Krasner, S. Total intensity of odor: A new method to evaluate odors. *J. Am. Water Works Assoc.* 2017, *109*, E42–E48. [CrossRef]
- Knaapila, A.; Raittola, A.; Sandell, M.; Yang, B. Self-ratings of olfactory performance and odor annoyance are associated with the affective impact of odor, but not with smell test results. *Perception* 2017, 46, 352–365. [CrossRef]
- 6. Zhang, X. Comparison and research progress on determination and evaluation methods of odor monitoring. *Chin. Environ. Prot. Ind.* **2018**, *11*, 63–65.
- 7. Nicell, J.A. Assessment and regulation of odour impact. Atmos. Environ. 2009, 43, 196–206. [CrossRef]
- 8. Vasile, A.; Danciulescu, V.; Gheorghita, T.; Kim, L.; Dediu, V. Environmental Impact Assessment of Odours from Solid Waste Landfills Located in Highly Urbanized Areas. *Rev. Chim.* **2017**, *68*, 1749–1751.
- 9. Sucker, K.; Both, R.; Bischoff, M.; Guski, R.; Winneke, G. Odor freguency and odor annoyance Part II: Dose-Response association and their modification by hedonic tone. *Int. Arch. Occup. Environ. Health* **2008**, *81*, 683–694. [CrossRef]
- Both, R.; Koe, L.C.C.; Liang, D.T.; Sucker, K.; Winneke, G.; Stuetz, R.M.; Koch, E. Odour intensity and hedonic tone: Important parameters to describe odour annoyance to residents? *Water Sci. Technol.* 2004, *50*, 83–92. [CrossRef]
- 11. Invernizzi, M.; Capelli, L.; Sironi, S. Proposal of odor nuisance index as urban planning tool. *Chem. Senses* **2017**, *42*, 105–110. [CrossRef]
- 12. Keck, M.; Mager, K.; Weber, K.; Keller, M.; Frei, M.; Steiner, B.; Schrade, S. Odour impact from farms with animal husbandry and biogas facilities. *Sci. Total Environ.* **2018**, *645*, 1432–1443. [CrossRef]
- 13. Bull, M.A.; Fromant, M.L. The performance of numerical odour assessment for the prediction of odour complaints from wastewater treatment works. *J. Chart. Inst. Water E* **2014**, *28*, 316–322. [CrossRef]
- 14. Winneke, G.; Kastka, J. Comparison of odor annoyance-data from different industrial sources: Problems and implications. In *Environmental Annoyance. Characterization, Measurement and Control*; Koelega, H.S., Ed.; Elsevier: Amsterdam, The Netherland, 1987; pp. 129–137.
- 15. Andersen, B.V.; Brockhoff, P.B.; Hylding, G. The importance of liking of appearance, -odour, -taste and -texture in the evaluation of overall liking. A comparison with the evaluation of sensory satisfaction. *Food Qual. Prefer.* **2019**, *71*, 228–232. [CrossRef]
- 16. Miedema, H.M.E.; Walpot, J.I.; Vos, H.; Steunenberg, C.F. Exposure-annoyance relationships for odour from industrial sources. *Atmos. Environ.* **2000**, *34*, 2927–2936. [CrossRef]
- 17. Chrea, C.; Valentin, D.; Sulmont-Rossé, C. Culture and odor categorization: Agreement between cultures depends upon the odors. *Food Qual. Prefer.* **2004**, *15*, 669–679. [CrossRef]
- 18. Qu, G.; Edeogu, I.E.; Feddes, J.J.R. Odor index: An integration of odor parameters for swine slurry odors. *Trans. ASABE* **2010**, *53*, 219–223. [CrossRef]
- Perry, E.; Al Ain, S.; Raineki, C.; Sullivan, M.; Wilson, A. Development of odor hedonics: Experience-dependent ontogeny of circuits supporting maternal and predator odor responses in rats. *J. Neurosci.* 2016, *36*, 6634–6650. [CrossRef]
- 20. Wang, J.; Shen, S.; Zhang, J.J. The college students' odour words classification-The discussion based on the semantic similarity and consciousness similarity. *Acta Psychol. Sin.* **2011**, *10*, 1124–1137.

- 21. Sironi, S.; Capelli, L.; Detoni, L. Methodology for assessing the performance of urine absorbing aids in controlling malodour release. *Int. J. Cometic Sci.* **2013**, *35*, 321–328. [CrossRef]
- 22. Yael, L.; Pola, O.; Uzi, R.; Rima, B.; Aviva, H.; Ibrahim, S.; Yael, A. A screening tool for selection of field odor assessors. *J. Air Waste Manag. Assoc.* 2011, *61*, 1353–1360.
- 23. Nimmermark, S. Characterization of Odor from Livestock and Poultry Operations by the Hedonic Tone. In Proceedings of the American Society of Agricultural and Biological Engineers annual international meeting, Portland, OR, USA, 9–12 July 2006; Volume 8 of 12, pp. 5296–5306.
- 24. Griffiths, K.D. Disentangling the frequency and intensity dimensions of nuisance odour, and implications for jurisdictional odour impact criteria. *Atmos. Environ.* **2016**, *90*, 125–132. [CrossRef]
- 25. Nimmermark, S. Influence of odour concentration and individual odour thresholds on the hedonic tone of odour from animal production. *Biosyst. Eng.* **2011**, *108*, 211–219. [CrossRef]
- 26. Huang, D.D.; Guo, H.P. Relationships between odor properties and determination of odor concentration limits in odor impact criteria for poultry and dairy barns. *Sci. Total Environ.* **2018**, *630*, 1484–1491. [CrossRef]
- 27. GB/T 14675-1993. *Air Quality–Determination of Odor–Triangle Odor Bag Method;* China Environmental Press: Beijing, China, 1993.
- 28. Chang, F.L.; Heinemann, P.H. Prediction of human assessments of dairy odor utilizing a fast gas chromatograph and neural networks. *Comput. Electron. Agric.* **2019**, *157*, 541–548. [CrossRef]
- 29. Opatz, O.; Soiffer, A.; Hummel, T. Are there gender related differences of responses to repetitive intranasal chemosensory stimuli? *Chem. Sci.* 2000, *25*, 797–805.
- Dalton, P.; Doolittle, N.; Berslin, P.A.S. Gender-specific induction of enhanced sensitivity to odours. *Nat. Neurosci.* 2002, 5, 199–200. [CrossRef]



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).